


# AHZ Allgemeine Homöopathische Zeitung


gegründet 1832

Wissenschaftliche und praktische Homöopathie

Abstracts of the 60th Congress of the Liga Medicorum Homoeopathica Internationalis - Berlin, May 4-7, 2005  
Abstractband des 60. Kongresses der Liga Medicorum Homoeopathica Internationalis - Berlin, 4.-7. Mai 2005



## HOEWAERTELICA INTERNATIONALIS



Small text at the bottom right of the page, likely a contact or publication information.

### Scope and limitation of homeopathy in rheumatoid arthritis

Horváth Katalin, Csontos Balázs, Székely Zoltán

5840 patients that fulfil the more than 25 million DM patients, this will be around 12 million by the end of 2005, making it the disease of the first choice. Keeping this in mind, a study was made to determine the place of homeopathy in the treatment of RA.

**Purpose:** To study the prevalence of RAHHS from insulin-dependent DM in different socio-economic groups and different races. Etiologies was also given an analysis of the risk factors for development of RAHHS and the efficacy of homeopathy as a treatment with various pathologies including DM and metabolic disorders.

**Methods:** 431 patients were taken into a 4000 two groups from all categories; they were treated to determine diaphoretic medicines.

**Group 1:** 215 patients who were into a single diet and exercise.  
**Group 2:** 215 patients who were given a single diet and exercise course.

**Results:** In Group 1, 14 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 2, 10 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 3, 170 patients responded to be insipidic medicines. 60 developed complications and were referred to allopathic doctors.

**Conclusion:** The constitutional medicine selected on the basis of the history of symptoms selected after proper case taking are effective for RAHHS. Some authors mention insulin DM also. Some homeopathic medicines have shown efficacy to control in the DM in different stages.

**Keywords:** Rheumatoid arthritis, diaphoretic medicines, insulin dependent diabetes mellitus, homeopathy.

### Homeopathic treatment of rheumatoid arthritis

Horváth Katalin, Csontos Balázs, Székely Zoltán

5840 patients that fulfil the more than 25 million DM patients, this will be around 12 million by the end of 2005, making it the disease of the first choice. Keeping this in mind, a study was made to determine the place of homeopathy in the treatment of RA.

**Purpose:** To study the prevalence of RAHHS from insulin-dependent DM in different socio-economic groups and different races. Etiologies was also given an analysis of the risk factors for development of RAHHS and the efficacy of homeopathy as a treatment with various pathologies including DM and metabolic disorders.

**Methods:** 431 patients were taken into a 4000 two groups from all categories; they were treated to determine diaphoretic medicines.

**Group 1:** 215 patients who were into a single diet and exercise.  
**Group 2:** 215 patients who were given a single diet and exercise course.

**Results:** In Group 1, 14 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 2, 10 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 3, 170 patients responded to be insipidic medicines. 60 developed complications and were referred to allopathic doctors.

**Conclusion:** The constitutional medicine selected on the basis of the history of symptoms selected after proper case taking are effective for RAHHS. Some authors mention insulin DM also. Some homeopathic medicines have shown efficacy to control in the DM in different stages.

**Keywords:** Rheumatoid arthritis, diaphoretic medicines, insulin dependent diabetes mellitus, homeopathy.

### Homeopathic treatment of rheumatoid arthritis

Horváth Katalin, Csontos Balázs, Székely Zoltán

5840 patients that fulfil the more than 25 million DM patients, this will be around 12 million by the end of 2005, making it the disease of the first choice. Keeping this in mind, a study was made to determine the place of homeopathy in the treatment of RA.

**Purpose:** To study the prevalence of RAHHS from insulin-dependent DM in different socio-economic groups and different races. Etiologies was also given an analysis of the risk factors for development of RAHHS and the efficacy of homeopathy as a treatment with various pathologies including DM and metabolic disorders.

**Methods:** 431 patients were taken into a 4000 two groups from all categories; they were treated to determine diaphoretic medicines.

**Group 1:** 215 patients who were into a single diet and exercise.  
**Group 2:** 215 patients who were given a single diet and exercise course.

**Results:** In Group 1, 14 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 2, 10 patients were only controlled by proper diet and exercise and 10 had relapsed. In Group 3, 170 patients responded to be insipidic medicines. 60 developed complications and were referred to allopathic doctors.

**Conclusion:** The constitutional medicine selected on the basis of the history of symptoms selected after proper case taking are effective for RAHHS. Some authors mention insulin DM also. Some homeopathic medicines have shown efficacy to control in the DM in different stages.

**Keywords:** Rheumatoid arthritis, diaphoretic medicines, insulin dependent diabetes mellitus, homeopathy.